
















FORMIVAR 60%	CH-Kasten Ruche Suisse (Bürki) Arnia Svizzera	Nuclei Mini Plus Warré	Zander DNM Langstroth Segeberger National, Simplex Smith, 2× Warré	Dadant 2× Simplex AZ Alberti- Žnideršič Commercial Deep National	2× Zander 2× DNM 2× Segeberger 2× Langstroth 4× Warré
	20–25 ml/d	20–25 ml/d	20–25 ml/d	25–30 ml/d	30–50 ml/d
25–30°C	1/4 	1/8 	2×7/8 	2×1 	2×1 
	1.100 ml	2.150 ml	1.150 ml 2.150 ml	1.200 ml	2.200 ml
20–25°C	3/8 	1/4 	2×1 	2×1 	2×1 
	1.100 ml	2.150 ml	1.150 ml 2.150 ml	1.200 ml	2.200 ml
15–20°C	2×3/4 	2×1/2 	2×1 	2×1 	2×1 
	1.100 ml	2.150 ml	1.150 ml 2.150 ml	1.200 ml	2.200 ml
















d Tag, day, jour, giorno, dag

1. Erste Behandlung, first treatment, premier traitement, primo trattamento, eerste behandeling
2. Zweite Behandlung, second treatment, deuxième traitement, secondo trattamento, tweede behandeling

Verwendung von Formivar 85% wird empfohlen.
Use of Formivar 85% is recommended.
L'utilisation de Formivar 85% est recommandée.
L'uso del Formivar 85% è consigliato.
Het gebruik van Formivar 85% wordt aanbevolen.



Verwenden Sie 2 Dochtpapiere pro Behandlung.
Use 2 paper-wicks per treatment. Utilisez 2 papiers par traitement. Utilizzare 2 carte per trattamento.
Gebruik 2 verdampappapieren per behandeling.

FORMIVAR 85%	CH-Kasten Ruche Suisse (Bürki) Arnja Svizzera	Nuclei Mini Plus Warré	Zander DNM Langstroth Segeberger National, Simplex Smith, 2×Warré	Dadant 2×Simplex AZ Alberti- ŽnidarsíČ Commercial Deep National	2×Zander 2×DNM 2×Segeberger 2×Langstroth 4×Warré
	10–15 ml/d	10–15 ml/d	15–20 ml/d	15–20 ml/d	20–30 ml/d
25–30 °C	1/16 	1/16 	1/4 	5/8 	3/4 
	 1.50 ml	 2.100 ml	 1.75 ml 2.100 ml	 1.100 ml	 2.200 ml 
20–25 °C	1/8 	1/8 	3/8 	3/4 	7/8 
	 1.50 ml	 2.100 ml	 1.75 ml 2.100 ml	 1.100 ml	 2.200 ml
15–20 °C	1/4 	1/4 	1/2 	7/8 	1 
	 1.50 ml	 2.100 ml	 1.75 ml 2.100 ml	 1.100 ml	 2.200 ml

d Tag, day, jour, giorno, dag

1. Erste Behandlung, first treatment, premier traitement, primo trattamento, eerste behandeling
2. Zweite Behandlung, second treatment, deuxième traitement, secondo trattamento, tweede behandeling



Verwendung von Formivar 60% wird empfohlen.

Use of Formivar 60% is recommended.

L'utilisation de Formivar 60% est recommandée.

L'uso del Formivar 60% è consigliato.

Het gebruik van Formivar 60% wordt aanbevolen.